

SOUTH AUSTRALIA

JUNE 2025




## **Australia's Changing Climate**

#### What is climate change?

Increases in greenhouse gases lead to changes in the atmosphere and the physical and biogeochemical ocean environment.

Australia's sea surface temperature (SST) has warmed by over 1 °C since 1900¹. Ongoing ocean warming contributes to longer and more frequent marine heatwaves. As waters warm, they also become more stratified – altering circulation and currents and reducing nutrient availability in surface waters.

Increased atmospheric  $CO_2$  means more  $CO_2$  is absorbed by the ocean, leading to shifts in ocean chemistry and ocean acidification.





## What impacts are expected for the seafood sector?

The seafood we eat relies on healthy marine ecosystems. Ecosystem productivity can be impacted through changing nutrient cycles, increasing temperatures, and habitat loss. Shallow coastal and estuarine ecosystems are particularly vulnerable to climate change.

The impact of climate change to the seafood sector will vary by the degree of exposure to physical change and species vulnerability.

Image credit: Nicolas Job

#### How to use this report



Use **page 2** to find information on SST changes over the past year and forecasts for the next two months in your area.



Use **pages 3-4** to learn about potential future ocean conditions due to climate change. These projections are based on a 'business as usual' scenario and show two global warming levels: 1.5°C (expected around 2015-2034) and 2°C (expected around 2030-2049). The Paris Agreement aims to limit global warming to well below 2°C above pre-industrial levels.



Report your climate observations and go 'Fishing for Climate Answers'! Check out the <u>Sea Change Australia</u> website to log potential climate related observations in your area and submit your climate questions. Our team connects your questions with climate and fisheries experts who provide clear, practical answers.






SOUTH AUSTRALIA

**JUNE 2025** 



### **Observed SST anomalies in 2024-25**



Monthly SST anomalies from May 2024 to April 2025 (base period 1993-2016)<sup>3</sup>. Cooler waters were present in nearshore waters during May-July 2024, after which waters began to warm. Marine heatwave conditions began in September 2024, and have persisted through April 20254. Warm temperatures and calm conditions led to the development of a Harmful Algal Bloom in the region.


Similar SST anomaly maps from BOM can be accessed here.

Waters have been anomalously warm

in 2024-2025.

## Forecast Conditions: Jul-Aug 2025

Australia Sea Surface Temperature Anomaly Outlook



Issued: 11/06/25 Base Period: 1981-2018



See BOM for seasonal forecast outlooks for SST

Monthly SST anomalies are forecast to remain anomalously warm from July-August 2025<sup>5</sup>. Forecasts are updated regularly and can be accessed here.




SOUTH AUSTRALIA

JUNE 2025




## **Projected Change: Ocean Warming**

### Baseline



#### **Projected**






Waters are projected to warm, with longer-lasting marine heatwaves.


SSTs are projected to increase under climate change. The degree of warming is highest in the west.




Sea Surface Temperature








Marine heatwaves are projected to increase in duration, and this pattern is highest in the west. Under a 2°C global warming level (~2030-2049), this region is projected to experience marine heatwave conditions for much of the year (relative to the baseline).





-100 -50 0 50 100 150



0.5 -0.4 -0.3 -0.2 -0.1 0.0 0.1 0.2 0.3 0.4 0.5

projected to increase under climate change. The degree of warming is highest in shallow waters on the continental shelf.

**Bottom temperatures** are

Climate projections are from the Ocean Forecasting Australian Model (OFAM), which is a spatially-downscaled ocean model<sup>6.</sup> Projections are based on a 'business as usual' scenario (CMIP5) and show two global warming levels: 1.5°C (expected around 2015-2034) and 2°C (expected around 2030-2049) relative to a historical baseline (2001-2020).



SOUTH AUSTRALIA

**JUNE 2025** 



## **Projected Change: Ecosystem Productivity**


#### **Baseline**

## Historical baseline (2001-2020)

NPP (molC/m2/v)

1 2 3 4 5 6 7 8 9 10 11

#### **Projected**




0

Productivity is projected to decline but is uncertain.

Projections of **net primary productivity**, the base of the marine food web, show decreases in the west and increases in the east. Projections are highly uncertain<sup>6</sup>.

# **Exploitable Fish Biomass**

**Net Primary Productivity** 



Ecosystem models project changes in the biomass available to fisheries (animals 10g - 100kg) under a 2.0°C global warming level.

On average, there is a decline in biomass, but this is uncertain. Minimum and maximum biomass estimates show divergent patterns.

Biomass is estimated from 6 ecosystem models projected under CMIP6<sup>7</sup>. Ecosystem models integrate temperature, oxygen, salinity, acidity, nutrients, & primary productivity, among other variables.



## -40 -30 -20 -10 0 10 20 30 40 50

**Community Reflections and Feedback** 

Change in biomass (%)

#### We would love to hear from you!

We invite reflections from the community about how the fisheries and aquaculture sectors are experiencing climate change in your region. Your story could be featured in this space.

The Sea Change Australia team listens and adapts to the needs of the sector. We want to make sure these report cards are as useful and relevant as possible—so let us know what you think. Your feedback will help shape future updates throughout the project.

Contact enquiries@seachangeaustralia.org





